


THE SOURCE

www.thesourceimports.com

RIOJA D.O.Ca. and NAVARRA D.O.

Subzones, Terroirs & Geological Map

VERSION 2-2022



**BIZKAIKO
TXAKOLINA D.O.**

**RIOJA D.O.Ca.
RIOJA ALTA**

Dominant Rock Types:
Limestones, Marlstones

Other Rock Types Present:
Mudstones, Sandstones,
Conglomerates

Altitude Range:
400-715 meters

Rainfall (mm): 390

Temperature (°C):
-10/2/29/42

**ARABAKO
TXAKOLINA D.O.**

**RIOJA D.O.Ca.
RIOJA ALAVESA**

Dominant Rock Types:
Limestones, Marlstones

Other Rock Types Present:
Mudstones, Sandstones, Conglomerates

Altitude Range:
400-750 meters

Rainfall (mm): 445

Temperature (°C):
-12/1/28/42

**NAVARRA D.O.
TIERRA ESTELLA**

Dominant Rock Types:
Highly variable

Other Rock Types Present:
Limestones, Marlstones, Mudstones,
Sandstones, Conglomerates, Rock Gypsum

Altitude Range:
350-700 meters

Rainfall (mm): 454-700

Temperature (°C): -11/2/29/42

**NAVARRA D.O.
VALDIZARBE**

Dominant Rock Types:
Highly variable

Other Rock Types Present:
Limestones, Marlstones, Mudstones, Sandstones,
Conglomerates, Rock Gypsum

Altitude Range:
350-500 meters

Rainfall (mm): 544- 807

Temperature (°C): -12/1/29/41

**NAVARRA D.O.
BAJA MONTAÑA**

Dominant Rock Types:
Highly variable

Other Rock Types Present:
Limestones, Marlstones, Mudstones,
Sandstones, Conglomerates

Altitude Range:
350-650 meters

Rainfall (mm): 470-760

Temperature (°C): -11/1/29/40

★ **Aseginolaza
& Leunda**

**NAVARRA D.O.
RIBERA ALTA**

Dominant Rock Types:
Highly variable

Other Rock Types Present:
Limestones, Marlstones, Mudstones,
Sandstones, Conglomerates,
Rock Gypsum

Altitude Range:
350-550 meters

Rainfall (mm): 349-507

Temperature (°C): -11/2/30/39

**NAVARRA D.O.
RIBERA BAJA**

Dominant Rock Types:
Highly variable

Other Rock Types Present:
Limestones, Marlstones, Mudstones,
Sandstones, Conglomerates

Altitude Range:
300-400 meters

Rainfall (mm): 361-384

Temperature (°C): -10/3/32/43

GRAPES

Reds

Tempranillo: M-HA, HT, DC
Garnacha: M-HA, HT, M-DC
Mazuelo (Cariñena) - M-HA, HT, DC
Graciano: HA, HT, DC
Maturana Tinta: HA, HT, DC

Whites

Viura (Macabeo): L-MA
Maturana Blanca: HA
Garnacha Blanca: L-MA
Tempranillo Blanco: M-HA
Turruntés: HA
Verdejo: HA
Alarije (Malvasía Riojana): M-HA
Moscatel de Grano Menudo: MA

Key

HA: High Acidity
MA: Medium Acidity
LA: Low Acidity
HT: High Tannin
MT: Medium Tannin
LT: Low Tannin
DC: Dark Color
MC: Medium Color
LC: Light Color

Madrid ~200km



Arizcuren ☆

**RIOJA D.O.Ca.
RIOJA ORIENTAL**

Dominant Rock Types:
High variability

Other Rock Types Present:
Mudstones, Sandstones,
Conglomerates,
Limestones, Marlstones

Altitude Range:
300-800 meters

Rainfall (mm): 350

Temperature (°C): -12/1/29/43

**CAMPO DE
BORJA D.O.**

This document is the work of Ted Vance, The Source Imports, and Ivan Rodríguez, MSc Geology.
The geological map is taken from "Mapa Geológico de España y Portugal (2015)".

The wine sub-region outlines were based on the official map of each respective appellation website.

It should be noted that the different colors on the geological map represent rocks from different geological time periods, and not specifically bedrock types.



30 km


18.64 miles

THE SOURCE


www.thesourceimports.com
GEOLOGICAL KEY

VERSION 2021-1


IBERIAN CONTINENTAL BASINS




IBERIAN VARISCAN MASSIF



IBERIAN MASSIF ALLOCHTHON UNITS



VARISCAN IGNEOUS ROCKS



Igneous Intrusive	(F) Felsic	(I) Intermediate	(M) Mafic
Igneous Extrusive	(F) Felsic	(I) Intermediate	(M) Mafic
Sedimentary	(C) Calcareous	(S) Siliciclastic	(O) Organic
Metamorphic	(L) Low-grade	(M) Medium-grade	(H) High-grade
	(N) Non-foliated (any metamorphism grade)		

Mya: Million years ago.

Massif: Compact group of mountains connected forming an independent portion of land.

Continental Basin: Topographically low inland area where sediments can accumulate.

Allochthon Unit: A large block of rock moved tectonically from where it was formed.

Variscan: related to the Variscan orogeny. An ancient chain of mountains which was the result of the formation of Pangea.

Not present in this area of the Iberian Peninsula:

1, 2, 4-7, 11, 14, 18-20, 38-42, 46, 48-52, 61-62, 65-70 71-75, 77-82, 84, 88-160, 163, 167-170, 172

THE SOURCE

TERROIR MAP SUPPORT MATERIAL


This series of maps is intended to provide an understandable yet concise collection of data that provides useful context for the world's curious wine tasters. While science has already produced a great deal of research on the nature of wine, there is still so much more work to be done.

Unlike many other terroir imprints, wine's relationship to rock and its mineral composition remains scientifically ambiguous, despite some observable patterns by people who have committed their lives to the subject. Within the science community there is a great deal of skepticism about this relationship, however history demonstrates that an open embrace of possibilities allows science to reach beyond its current limits.

The technical elements of this work has been reviewed by researchers from the Department of Marine Geoscience and Planning Territory of the University of Vigo, Spain.

IGNEOUS ROCK

- Rock formed from cooling of magma.
- Intrusive/Plutonic: cooled slowly below the surface.
- Extrusive/Volcanic: cooled quickly above the surface.
- The Mineral crystal size of igneous rocks depends on the rate the magma cools. Slow cooling results in larger crystals than those that cool faster. Crystal size affects the soil grain size and has implications with soil water retention, which in turn influences a resulting wine's characteristics.
- Felsic, Intermediate and Mafic (also, Ultramafic) charts igneous rock mineral composition (see diagram 1).
- Intrusive and extrusive rocks from the same magma chamber share the same mineral makeup.
- Igneous rock pH levels are related to the composition classification. Felsic are more acidic, while the pH increases as we move to Mafic and Ultramafic.



METAMORPHIC ROCK

- Rock altered (metamorphosed) from a pre-existent rock (protolith) by extreme conditions of pressure and/or temperature.
- Depending on how much these conditions affected the rock, they could be classified as low, medium or high-grade metamorphic rock.
- The main divisions of metamorphic rock are based on appearance and classified as either foliated or non-foliated and how pressure/temperature affected the preexisting rock.
- Each classification and location of metamorphic rock has a unique mineral composition, which leads to variance in modes of decomposition.
- Metamorphic rock pH levels mainly depends on the acidic/alkaline nature of the protolith.

ROCK NAME	TEXTURE	GRAIN SIZE	PARENT ROCK (PROTOLITH)
SLATE	FOLIATED	FINE	SHALE, MUSTONE OR SILTSTONE
SCHIST		MEDIUM TO COARSE	SLATE OR IGNEOUS ROCK
GNEISS		MEDIUM TO COARSE	SCHIST OR IGNEOUS ROCK
MARBLE	NON-FOLIATED	MEDIUM TO COARSE	CALCAREOUS ROCK
QUARTZITE		MEDIUM TO COARSE	QUARTZ SANDSTONE
ANTHRACITE		FINE	COAL

Modified from 2006 Pearson Prentice Hall, Inc.

SEDIMENTARY ROCK

- Rock formed from the accumulation of sediments, compaction and cementation (lithification).
- Three main types:
 - Siliciclastic: fragments of igneous and metamorphic rocks, mainly silicates.
 - Calcareous: mainly fragments of carbonate-rich organisms.
 - Organic-rich: containing significant amounts of organic carbon of biological origin.
- In general, pH levels in siliciclastic are acidic while calcareous are alkaline.

SILICICLASTIC			CHEMICAL	
GRAIN SIZE	SEDIMENT	ROCK	COMPOSITION	ROCK
COARSE (>2 MM)	ROUNDED GRAVEL	CONGLOMERATE	CALCITE (CaCO ₃)	LIMESTONE
	ANGULAR GRAVEL	BRECCIA	QUARTZ (SiO ₂)	CHERT / FLINT
MEDIUM (2 TO 0.062 MM)	SAND	SANDSTONE	GYPHUM (CaSO ₄ ·2H ₂ O)	GYPHUM
FINE (0.062 TO 0.004 MM)	SILT	SILTSTONE	HALITE (NaCl)	HALITE
VERY FINE (<0.004 MM)	CLAY	SHALE	PLANT FRAGMENTS (C+others)	COAL

THE SOURCE

WEATHERING + TOPSOIL COMPOSITION

- A vineyard's soil depth, grain and mineralogy influence a resulting wine's organoleptic characteristics.
- Erosion depends on mineral composition and degree of exposure within the environment.
- Rocks erode either by physical weathering (fragmentation) and/or chemical weathering (mineral dissolution).
- Physical weathering produces coarser soils. Chemical weathering produces finer soils.
- Inorganic components of soil are bedrock-derived and/or unconsolidated (transported) material.
- Soil tends to be composed of a mix of grains, which are broadly classified (small to large) as clay, silt, sand, or gravel.

CLAY

- Clay has more significant implications compared to other soil grains due to its increased water retention capacity and nutrient/mineral supply—a consequence of its smaller size and more complex composition.
- Clay has a higher CEC (Cation Exchange Capacity) than other soil grains, and each clay type has a different CEC and water retentive capacity. CEC level is directly related to fertility: e.g., low CEC implies low fertility. (Cation: An atom with positive electrical charge which allows them to be highly reactive with other atoms.)
- Four *main* clay mineral types:
 - Kaolinite has the simplest chemical composition, largest soil grain, relatively low CEC, and low water retention; it is referred to as a *non-swelling clay*. It's commonly a byproduct of weathered feldspar and muscovite from felsic rocks (rhyolite and granite), and allophane from recent volcanic rocks (tephra).
 - Illite is a potassium aluminum-rich clay with intermediate CEC and intermediate water retention. It is commonly a byproduct of weathered felsic igneous or high-grade metamorphic rocks.
 - Vermiculite is an iron and magnesium rich clay, with high CEC and intermediate water retention. It is commonly a byproduct of weathered biotite, chlorite and amphiboles (usually from volcanic rock).
 - Smectite is fine-grained with high CEC and high water retention. It is commonly a product of weathered basalts and other volcanic rocks. In sediments, these clay minerals have various origins and are most often reworked from soils where clay formed under warm, humid climates. It also forms in marine environments at the expense of volcanic glass or formed after slow deposition of sediments.

TEMPERATURE AND PRECIPITATION

- The Köppen climate classification (categorized as Oceanic, Continental, Mediterranean, etc.) is useful on a global scale, but concerning wine regions it can be vague and confusing (e.g. Galicia and all of Portugal are classified as Mediterranean). We have opted instead to use regional temperature and precipitation data.
- While generalities with precipitation and temperature are useful, geographical characteristics like altitude, slope, exposure, wind currents, and proximity to hills, mountains, water (rivers, oceans, seas, lakes) influence temperature, precipitation and humidity are specific to each terroir.
- Temperatures are arranged as lowest/average low/average high/highest (e.g. -4/0/28/33). These numbers better illustrate a region's climate than broad categorizations. It may also help with general concepts. Examples:
 - -7/1/30/43 from Portugal's Planalto Mirandês (PM) compared to -3, 7, 25, 38 from Spain's Val do Salnés (VdS) clearly demonstrates that winter lows and summer highs in PM are more extreme than the VdS which is more moderate on both ends.
 - PM vineyards may be affected by frost because temperatures below zero may happen after vegetative cycles begin, while in VdS vineyards are not affected (yet) because the temperatures may only drop below zero in the winter while vines are dormant.
 - Because of VdS's moderate temperature plant life cycles in Spring generally start sooner than PM. There are exceptions for specific grape varieties with earlier or later vegetative cycles.
- Seasonal temperature average ranges (the middle numbers, e.g. -4/0/28/33) were gathered from weatherspark.com in 2021 and are based on data from 1980 to 2016. Regional winegrowers have supplied extreme cold and hot temperatures witnessed (-4/0/28/33) up to 2021, within specific vineyard zones.
- Temperatures are listed in Celsius (°C) and precipitation in millimeters (mm). (To convert °C to Fahrenheit (°F) multiply °C by 1.8 and add 32. To convert mms to inches multiply mms by 0.0393701.)
- Precipitation levels (listed on maps as an annual average) demonstrate the overall climatic conditions and landscape setting and prompt many considerations. A few examples:
 - Low precipitation: low humidity and low mildew pressure, fewer vineyard treatments needed against mildew and thereby overall easier for ecological and organic viticulture.
 - High precipitation: high humidity and high mildew pressure, so more vineyard treatments are needed against mildew and it's more difficult for ecological and organic viticulture.